compounds that may be used in the *methods* of the present invention, as described in this section of the specification, does not contain the provisos set forth in the section of the specification on pages 20 to 26, that is directed specifically to *compounds*. The provisos were included in the latter section, because certain species covered by the generic disclosure had been described previously. However, the use of such compounds in the methods defined by the pending claims was neither taught nor suggested in that art. Accordingly, the genus of compounds suitable in the claimed methods, as described at pages 14 to 20, is broader than the genus of novel compounds, described at pages 20 to 26.

Applicants respectfully submit that the full scope of the invention, as defined by the pending claims, is described in the specification, as required by 35 U.S.C. § 112, first paragraph. Accordingly, Applicants respectfully request that the rejection be withdrawn.

Copies of the references cited in the IDS submitted on August 16, 2001 are enclosed herewith. Consideration of the references is respectfully requested.

Applicants acknowledge the non-statutory double patenting rejection over Claims 28 and 29 of co-owned U.S. Patent No. 6,191,131. Applicants propose to file a terminal disclaimer to overcome this rejection upon receiving an indication from the Examiner that the claims otherwise define allowable subject matter.

If there are any additional issues, or if the Examiner wishes to discuss this application further, please telephone the Applicants' undersigned representative at the telephone number below. Applicants note, however, that all written correspondence for this application should continue to be sent to:

Bristol-Myers Squibb Pharma Company
Patent Department
PO Box 4000
Route 206 and Provinceline Road
Princeton, NJ 08453-4000

Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "**Version with markings to show changes made.**"

Date: 2/21/03

S. Maurice Valla
Registration No. 43,966

Woodcock Washburn LLP
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone: (215) 568-3100
Facsimile: (215) 568-3439

VERSION WITH MARKINGS TO SHOW CHANGES MADE

Claim 1 has been re-typed, as follows:

1. A method of treating a disorder induced or facilitated by CRF in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of Formulae (1) or (2):



(1)



(2)

isomers thereof, stereoisomeric forms thereof, mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt forms thereof, wherein:

Z is CR^2;

Ar is selected from phenyl, naphthyl, pyridyl, pyrimidinyl, triazinyl, furanyl, thienyl, benzothienyl, benzofuranyl, 2,3-dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, indanyl, 1,2-benzopyranyl, 3,4-dihydro-1,2-benzopyranyl, tetralinyl, each Ar optionally substituted with 1 to 5 R^4 groups and each Ar is attached to an unsaturated carbon atom;

R^1 is independently selected at each occurrence from H, C_1-C_4 alkyl, C_2-C_4 alkenyl, C_2-C_4 alkynyl, halo, CN, C_1-C_4 haloalkyl, C_1-C_{12} hydroxyalkyl, C_2-C_{12} alkoxyalkyl, C_2-C_{10} cyanoalkyl, C_3-C_6 cycloalkyl, C_4-C_{10} cycloalkylalkyl, NR^9R^{10}, C_1-C_4 alkyl-NR^9R^{10}, NR^9COR^{10}, OR^{11}, SH or $S(O)_nR^{12}$;

R^2 is selected from H, C_1-C_4 alkyl, C_2-C_4 alkenyl, C_2-C_4 alkynyl, C_3-C_6 cycloalkyl,

C_4-C_{10} cycloalkylalkyl, C_1-C_4 hydroxyalkyl, halo, CN, -NR^6R^7, NR^9COR^{10}, -$NR^6S(O)_nR^7$, $S(O)_nNR^6R^7$, C_1-C_4 haloalkyl, -OR^7, SH or -$S(O)_nR^{12}$;

R^3 is selected from:

-H, SH, $S(O)_nR^{13}$, COR^7, CO_2R^7, $OC(O)R^{13}$, NR^8COR^7, $N(COR^7)_2$, $NR^8CONR^6R^7$, $NR^8CO_2R^{13}$, $N(OR^7)R^6$, $CONR^6R^7$, aryl, heteroaryl and heterocycle, or

-C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_3-C_8 cycloalkyl, C_5-C_8 cycloalkenyl, C_4-C_{12} cycloalkylalkyl or C_6-C_{10} cycloalkenylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C_1-C_6 alkyl, C_3-C_6 cycloalkyl, halo, C_1-C_4 haloalkyl, cyano, OR^{15}, SH, $S(O)_nR^{13}$, COR^{15}, CO_2R^{15}, $OC(O)R^{13}$, NR^8COR^{15}, $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{13}$, $NR^{16}R^{15}$, $CONR^{16}R^{15}$, aryl, heteroaryl and heterocyclyl;

R^4 is independently selected at each occurrence from: C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_3-C_6 cycloalkyl, C_4-C_{12} cycloalkylalkyl, NO_2, halo, CN, C_1-C_4 haloalkyl, NR^6R^7, NR^8COR^7, $NR^8CO_2R^7$, COR^7, OR^7, $CONR^6R^7$, $CO(NOR^9)R^7$, CO_2R^7, or $S(O)_nR^7$, where each such C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_3-C_6 cycloalkyl and C_4-C_{12} cycloalkylalkyl are optionally substituted with 1 to 3 substituents independently selected at each occurrence from C_1-C_4 alkyl, NO_2, halo, CN, NR^6R^7, NR^8COR^7, $NR^8CO_2R^7$, COR^7 OR^7, $CONR^6R^7$, CO_2R^7, $CO(NOR^9)R^7$, or $S(O)_nR^7$;

R^6, R^7, R^{6a} and R^{7a} are independently selected at each occurrence from:

-H,

-C_1-C_{10} alkyl, C_3-C_{10} alkenyl, C_3-C_{10} alkynyl, C_1-C_{10} haloalkyl with 1-10 halogens, C_2-C_8 alkoxyalkyl, C_3-C_6 cycloalkyl, C_4-C_{12} cycloalkylalkyl, C_5-C_{10} cycloalkenyl, or C_6-C_{14} cycloalkenylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C_1-C_6 alkyl, C_3-C_6 cycloalkyl, halo, C_1-C_4 haloalkyl, cyano, OR^{15}, SH, $S(O)_nR^{13}$, COR^{15}, CO_2R^{15}, $OC(O)R^{13}$, NR^8COR^{15}, $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{13}$, $NR^{16}R^{15}$, $CONR^{16}R^{15}$, aryl, heteroaryl or heterocyclyl,

-aryl, aryl(C_1-C_4 alkyl), heteroaryl, heteroaryl(C_1-C_4 alkyl), heterocyclyl or heterocyclyl(C_1-C_4 alkyl);

alternatively, NR^6R^7 and $NR^{6a}R^{7a}$ are independently piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine or thiomorpholine, each optionally substituted with 1-3 C_1-C_4 alkyl groups;

R^8 is independently selected at each occurrence from H or C_1-C_4 alkyl;

R^9 and R^{10} are independently selected at each occurrence from H, C_1-C_4 alkyl, or C_3-C_6 cycloalkyl;

R^{11} is selected from H, C_1-C_4 alkyl, C_1-C_4 haloalkyl, or C_3-C_6 cycloalkyl;

R^{12} is C_1-C_4 alkyl or C_1-C_4 haloalkyl;

R^{13} is selected from C_1-C_4 alkyl, C_1-C_4 haloalkyl, C_2-C_8 alkoxyalkyl, C_3-C_6 cycloalkyl, C_4-C_{12} cycloalkylalkyl, aryl, aryl(C_1-C_4 alkyl)-, heteroaryl or heteroaryl(C_1-C_4 alkyl)-;

R^{14} is selected from C_1-C_{10} alkyl, C_3-C_{10} alkenyl, C_3-C_{10} alkynyl, C_3-C_8 cycloalkyl, or C_4-C_{12} cycloalkylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C_1-C_6 alkyl, C_3-C_6 cycloalkyl, halo, C_1-C_4 haloalkyl, cyano, OR^{15}, SH, $S(O)_nR^{15}$, COR^{15}, CO_2R^{15}, $OC(O)R^{15}$, NR^8COR^{15}, $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{15}$, $NR^{16}R^{15}$, $CONR^{16}R^{15}$, and C_1-C_6 alkylthio, C_1-C_6 alkylsulfinyl and C_1-C_6 alkylsulfonyl;

R^{15} and R^{16} are independently selected at each occurrence from H, C_1-C_6 alkyl, C_3-C_{10} cycloalkyl, C_4-C_{16} cycloalkylalkyl, except that for $S(O)_nR^{15}$, R^{15} cannot be H;

aryl is phenyl or naphthyl, each optionally substituted with 1 to 5 substituents independently selected at each occurrence from C_1-C_6 alkyl, C_3-C_6 cycloalkyl, halo, C_1-C_4 haloalkyl, cyano, OR^{15}, SH, $S(O)_nR^{15}$, COR^{15}, CO_2R^{15}, $OC(O)R^{15}$, NR^8COR^{15}, $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{15}$, $NR^{16}R^{15}$, and $CONR^{16}R^{15}$;

heteroaryl is pyridyl, pyrimidinyl, triazinyl, furanyl, pyranyl, quinolinyl, isoquinolinyl,

thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, isoxazolyl, pyrazolyl, 2,3-dihydrobenzothienyl or 2,3-dihydrobenzofuranyl, each being optionally substituted with 1 to 5 substituents independently selected at each occurrence from C_1-C_6 alkyl, C_3-C_6 cycloalkyl, halo, C_1-C_4 haloalkyl, cyano, OR^{15}, SH, $S(O)_nR^{15}$, $-COR^{15}$, CO_2R^{15}, $OC(O)R^{15}$, NR^8COR^{15}, $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{15}$, $NR^{16}R^{15}$, and $CONR^{16}R^{15}$;

heterocyclyl is saturated or partially saturated heteroaryl, optionally substituted with 1 to 5 substituents independently selected at each occurrence from C_1-C_6 alkyl, C_3-C_6 cycloalkyl, halo, C_1-C_4 haloalkyl, cyano, OR^{15}, SH, $S(O)_nR^{15}$, COR^{15}, CO_2R^{15}, $OC(O)R^{15}$, NR^8COR^{15}, $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{15}$, $NR^{15}R^{16}$, and $CONR^{16}R^{15}$; and

n is independently at each occurrence 0, 1 or 2.